

2011 Fungicide Resistance Management Guidelines for Cucurbit Downy Mildew and Powdery Mildew Control in the Northeast United States

Fungicide	Active Ingredient(s)	FRAC Code*	Risk Rating**	Management Required***	Powdery Mildew	Downy Mildew	Fungicide Resistance Management Guidelines****
Kocide 3000 or OLF	fixed copper(s)	M1	L	N	+		FRAC code M fungicides are low risk, protectant fungicides. Use alone, or tank mix with high-risk fungicides to improve control
Microthiol or OLF	sulfur	M2	L	N	++		
Manzate, Dithane or OLF	EBDC	M3	L	N		++	
Bravo or OLF	chlorothalonil	M5	L	N	++	++	
Topsin M	thiophanate methyl	1	H ^R	Y	+		Select fungicides with at least ++ rating. Rotate among fungicides with different FRAC codes. Tank mix high risk fungicides with FRAC code M product if the product is not formulated with a FRAC code M fungicide. When resistance is qualitative (FRAC code 1 and 11 fungicides), resistant pathogen strains are completely insensitive and cannot be controlled with the fungicide. With quantitative resistance (FRAC Code 3 fungicides), pathogen strains exhibit range in fungicide sensitivity and efficacy depends on level of insensitivity. Better control can be obtained with highest label rates and shortest spray intervals. Specific notes: Newer code 3 fungicides expected to be as effective as older ones when applied at same rate of active ingredient. Revus ineffective on cucumber.
Rally	myclobutanil	3	M ^R	Y	++		
Procure	triflumizole	3	M ^R	Y	++		
Folicur	tebuconazole	3	M	Y	++		
Inspire Super	difenoconazole + cyprodinil	3 + 9	M	Y	++		
Ridomil Gold Copper	mefenoxam + copper	4 + M1	H ^R + L	Y		+	
Ridomil Gold Bravo	mefenoxam + chlorothalonil	4 + M5	H ^R + L	Y		+	
Quadris	azoxystrobin	11	H ^R	Y	+	+	
Cabrio	pyraclostrobin	11	H ^R	Y	+	+	
Flint	trifloxystrobin	11	H ^R	Y	+		
Reason	fenamidone	11	H	Y		+	
Pristine	pyraclostrobin + boscalid	11 + 7	H ^R	Y	+	+	
Tanos	fomoxadone + cymoxanil	11 + 27	M	Y		+	
Quintec	quinoxifen	13	H	Y	+++++		
Ranman	cyazofamid	21	M - H	Y		+++	
Gavel	zoxamide + mancozeb	22 + M3	M + L	Y		++	
Curzate	cymoxanil	27	L - M	Y		++	
Previcur Flex	propamocarb HCL	28	L - M	Y		+++	
Aliette	aluminum tris	33	L	Y		+	
Phosphonates	phosphorous acid salts	33	L	Y		+	
Forum	dimethomorph	40	L - M	Y		+	
Revus	mandipropamid	40	L - M	Y		+	
Presidio	fluopicolide	43	H	Y		++++	

Efficacy Ratings: + = poor (not recommended), ++ = poor to good, +++ = good, ++++ = very good, +++++ = excellent

* FRAC code: M = multi-site mode of action (MOA), numbered groups = fungicides with similar MOA

** Risk Ratings: L = low risk, M = moderate risk or H = high risk for fungicide resistance to develop

*** Risk management required according to fungicide label

**** See fungicide label for specific crops, rates and instructions on use

^R = resistance known; (+) control failures detected in the mid-Atlantic and Northeast regions.

Fungicides with the same color belong to the same FRAC code.

Trade or Brand Names Disclaimer: The trade or brand names given herein are supplied with the understanding that no discrimination is intended and no endorsement by the Rutgers Cooperative Extension is implied. Furthermore, in some instances the same compound may be sold under different names, which may vary as to label clearances.