

# Southern Region BMSB Update

2016 funding from SIMPC to develop a Southern BMSB working group; AL, GA, FL, SC, NC, TN



## Main goals

- Hold a southern region stakeholder meeting and develop a priority list for upcoming research; April
- Establish formal communications among working group members; Basecamp
- Update known hosts and distribution maps for the southern US; in progress

Monitoring: current distribution & expansion  
Overwintering: if and where  
Reproducing populations  
Establishment  
Host plants  
Crops/Damage

## Southern Brown Marmorated Stink Bug Working Group

The Southern Brown Marmorated Stink Bug (BMSB) working group was established to enhance the development and implementation of IPM for BMSB in the southern region of the US based on its biology, phenology, behavior, and ecology. Our group complements the efforts of the existing BMSB working group in the Northeast.

With BMSB moving into the Southern region where there is a great diversity and abundance of hosts, conventional and organic growers have growing concerns regarding the future impact of this invasive pest on cropping systems in the region. Indeed, previously established economic thresholds for stink bugs in southern field crops may need to be redefined for these crops. Further, homeowners and the pest control industry in this region have an increasing interest in IPM for BMSB. Also, there is great demand for educational materials that explain BMSB ecology in urban and agricultural landscapes.

## Partners



## Statistics

1,017 County Reports  
483 Point Reports

## Recent Reports

- Ana Gutierrez in Fayette County, Kentucky  
March 7, 2016
- kristine camerra in Davidson County, Tennessee  
March 1, 2016
- Gil Hearn in DeKalb County, Georgia  
February 21, 2016
- Steve Bunning in Albemarle County, Virginia  
October 23, 2015
- Patty Abernathy in Rutherford County, Tennessee  
October 22, 2015

With BMSB moving into the Southern region where there is a great diversity and abundance of hosts, conventional and organic growers have growing concerns regarding the future impact of this invasive pest on cropping systems in the region. Indeed, previously established economic thresholds for stink bugs in southern field crops may need to be redefined for these crops. Further, homeowners and the pest control industry in this region have an increasing interest in IPM for BMSB. Also, there is great demand for educational materials that explain BMSB ecology in urban and agricultural landscapes.

### Partners



United States Department of Agriculture

National Institute of Food and Agriculture

### Recent Reports

Ana Gutierrez in Fayette County, Kentucky  
March 7, 2016

kristine camerra in Davidson County, Tennessee  
March 1, 2016

Gil Hearn in DeKalb County, Georgia  
February 21, 2016

Steve Bunning in Albemarle County, Virginia  
October 23, 2015

Patty Abernathy in Rutherford County, Tennessee  
October 22, 2015

[More Reports](#)

### Educational Resources

- ✓ Stop BMSB
- ✓ Southern Region Brown Marmorated Stink Bug Working Group Flyer
- ✓ Southern Region Brown Marmorated Stink Bug Working Group
- ✓ Brown Marmorated Stink Bug in Georgia

# Identification

## Description

The Brown Marmorated Stink Bug (BMSB) adult is similar in appearance to several native stink bug species, but can be distinguished by several characteristics. Firstly, it has light-colored bands on the joints of the antenna and legs. Also, on its ventral side, its abdominal venter is white to ash grey and may include a black spot at the terminus of the abdomen. The thin beak extends between the hind legs and is always dark colored. On the dorsal side, adults have a brown or grey marbled appearance with blue-green metallic coloration in the depressions on the head, pronotum, and connexivum, which is mostly visible in sunlight. They also have light and dark colored spots around the lateral margins of the abdomen that are not covered by the wings when the insects are at rest. The leading edge of the shoulder is smooth and not pointed on the ends. Adults are approximately 2/3" long, and females are typically larger than males.



Brown/grey marbling on dorsal side; light and dark colored spots along margin of abdomen not covered by wings

# Images



Share Flag Fullscreen



Adult



Eggs



First instars



First instars



First instar and second instars

# Images



5443490

Share | Flag | Fullscreen

- First instars
- First instar and second instars
- Third instar with spiny protrusions
- Third instar**
- Third instar
- Fourth instars
- Fifth instar

Susan Ellis, Bugwood.org


Third instar

## Brown Stink Bug

*Euschistus servus*

No bands on the antennae

20



No metallic coloration in depressions

No bands on the legs

21



Female

22



Male

Ventral Surface green to yellow

23



Eggs

24



1<sup>st</sup> instars

25



2<sup>nd</sup> instar

26



3<sup>rd</sup> instar

27



4<sup>th</sup> instar

28



5<sup>th</sup> instar

### Other Closely Resembling Species:

29



*Brochymena quadripustulata*

30



*Euschistus tristigmus*

## Brown Marmorated Stink Bug

*Halyomorpha halys*

The brown marmorated stink bug is a native of China, Taiwan, Korea and Japan. This invasive insect pest has spread to the United States, Canada, and Europe.

It is a serious economic pest of orchard crops, including apple and peach, row crops such as corn and soybeans, and vegetable crops including sweet corn, pepper, eggplant, and tomato.

### Photo Credits

Gary Bernon, USDA, APHIS, Bugwood.org (14)

Susan Ellis, Bugwood.org (12,15,17-19)

Kristie Graham, USDA, ARS (1-11, 29)

Davis R. Lance, USDA, APHIS, PPQ, Bugwood.org (13)

Herb Pilcher, USDA, ARS (20-28,30)

Martin E. Rice, Pioneer Hi-Bred (16)

### Information

<http://www.sipmc.org/BMSB>

To report a BMSB sighting:

<http://www.EDDMapS.org/BMSB/Report>

Funding provided by USDA NIFA, under Agreement No. 2014-70006-22485 via Southern IPM Center Working Group Program (Project 999458)

## Southern Region Brown Marmorated Stink Bug Working Group



THE UNIVERSITY OF GEORGIA  
**COOPERATIVE  
EXTENSION**



United States Department of Agriculture


**Agricultural Research Service**

# ID flyer for extension agents, pest control operators, growers, researchers

## Brown Marmorated Stink Bug

*Halyomorpha halys*

The BMSB adult is similar in appearance to several native species. A few identifying differences between the BMSB and the brown stink bug (native) are shown below.



## Adults

Adults are approximately 2/3" long. In most cases the female is larger than the male.



Female



Male

## Eggs



Fresh



Close to hatching

## Eggs with Nymphs



Hatched egg mass with 1<sup>st</sup> instars



Hatched egg mass with 1<sup>st</sup> instars molting into 2<sup>nd</sup> instars

## Nymphs

Nymphs have five instars (immature stages) that range from 1/16" to 1/2" in length. They move quickly – faster than the native species.



3<sup>rd</sup> instar



4<sup>th</sup> instar



Older instars begin showing white bands on their legs

5<sup>th</sup> instar wings developing



Younger instars have serrated (spikes) edges along the pronotum



## Got Brown Marmorated Stink Bugs? Where did you see it?



In a home, vehicle or structure



In a orchard, field, or crop



Can choose type of report

EDDMapS

www.eddmaps.org/bmsb/report/home.cfm

Home Report Sightings Maps Identification Resources Contact sign out

Red fields are required.

**Pest (?) :**  
Halyomorpha halys (brown marmorated stink bug)

**Observation Date: (?) :**  
06/14/2016

**Life Stage(s) Observed:**  
 Adult  Nymph  Egg

**Incidence (?) :**  
10 %

**Structure type (?) :**  
brick wall

**State:**

**County:**

**Latitude (?) :**

**Longitude (?) :**

**Host plant species:**

**Number Observed:**

**Severity (?) :**  %

**Substrate (?) :**



Map Satellite

12:10 PM  
6/14/2016

Nik Wiman working with Joe LaForest to develop a more detailed report.

EDDMapS

www.eddmaps.org/bmsb/report/crop.cfm

Red fields are required.

**Pest (?) :**  
Halyomorpha halys (brown marmorated stink bug)

**Observation Date: (?) :**  
06/14/2016

**Life Stage(s) Observed:**  
 Adult  Nymph  Egg

**Incidence (?) :**  
10 %

**Prevalence within stink bug community:**  
5 %

**State:**  
Georgia

**County:**  
Peach County

**Latitude (?) :**  
Must be expressed in Decimal Degrees (XX.XXXX), and DATUM NAD83/WGS84.

**Longitude (?) :**  
Must be expressed in Decimal Degrees (XX.XXXX), and DATUM NAD83/WGS84.

lat/long conversion tools place marker at position clear map


**Location Description/Nearest Address:**

**Host plant species:**  
black cherry

**Habitat (?) :**  
Edge: Field/forest

**Sex:**  
 Unknown  Female  Male

**Severity (?) :**  
%




Map Satellite

12:14 PM  
6/14/2016

Reproductive populations?





State verifiers

StopBMSB can this if want. Contact Joe LaForest with UGA in Tifton, GA.


# Using Pheromone-baited Pyramid Traps to Monitor BMSB

For traps in the southern region, an insect-collecting device made from an aerated clear plastic jar with several air vents is seated atop a black corrugated plastic pyramid base (Great Lakes IPM, Inc.). The insertion of three eyelets in the lid of the insect-collecting device allows adult stink bug parasitoids, but not stink bug adults, to escape.

AgBio, Inc. combo lure: BMSB aggregation pheromone + *P. stali* aggregation pheromone




*Trichopoda* spp. adult



# BMSB In Florida: Detected, but not established

- No evidence of a sustained, reproducing population in Florida.
- 36 detections consisting of 69 specimens.
- Additional detections usually associated with travelers returning from states with BMSB – mostly at Florida's agricultural interdiction stations or on RVs.



Figures adapted from [www.stopbmsb.org](http://www.stopbmsb.org)

# BMSB in Florida Peach

- 28 yellow pyramid traps baited with BMSB pheromone lure deployed at 5 locations
- 2 Adult BMSB recovered from orchard in Lake County (note: there was a traveler from northern states that could account for these captures)
- No additional finds at positive location in 60 day period following detection
- No nymphs or signs of reproduction
- At this time do not anticipate that BMSB will be a problem in Florida peach



# Georgia BMSB Team

Joe LaForest

Ted Cottrell

Glynn Tillman

Michael Toews

David Buntin

Rick Hoebeke

Dan Suiter

Lisa Ames

Phillip Roberts


Elizabeth Moss

Ash Sial

Whitney Hadden


# Known Distribution of BMSB in Georgia

- 2011-2016: Urban pest management professionals reporting overwintering aggregations in the Atlanta Metro area; established populations; recently egg mass found on apple in Cherokee county
- 2014: reproducing BMSB populations in cotton, pecan, catulpa, and ornamental hibiscus
- 2014: adults in cotton
- Jan. 2015: authorities at the Port of Savannah disclosed detection of BMSB in international cargo shipped out of Ga.
- 2015: BMSB on apples in Gilmer County in the Blue Ridge Mountain region
- 2015: reproducing BMSB populations in soybean in Pike County in the Piedmont region
- 2015: reproducing BMSB populations in peach in Peach County in the Coastal Plain region
- Adults detected
- I-75 Hitchhiker



Status: limited establishment, major nuisance pest in homes, reproductive populations in some agricultural crops

# BMSB invades the coastal plain in 2015




**Byron, GA** - late instars found on peach Aug. 2015.

- trapping during fall 2015 revealed adults at the SEFTNRL.

Ted Cottrell, USDA, ARS

# BMSB adults detected in Byron, GA in Spring 2016




Traps are near peach/pecan/woods. BMSB overwintered successfully at Byron – adults captured in fall of 2015, detected in homes during the winter, captured adults in spring of 2016. Established in this area – nymphs found late season in 2015 and overwintering adults captured the following spring.


Ted Cottrell, USDA, ARS

# First Detection of BMSB in Peach Away from Byron, GA

This past Friday (June 10), 2 BMSB adults were captured in traps near a commercial peach orchard about 12 miles southwest of Byron. All things considered, their numbers are low compared with the brown stink bug. This is not far from Fort Valley where Hwy 49 and 341 are major thoroughfares. So it's not farfetched that BMSB was brought in from other areas via vehicles.



# BMSB at UGA Bledsoe Row Crop Farm near Griffin, GA



- nymphs and adults in soybean 2015
- some OW adults observed in March in Griffin, GA
- OW adults captured in spring of 2016
- established in the county


2016 trap captures:

5/16 - 5 adults  
5/23 - 3 adults  
5/30 - 3 adults  
6/6 - 3 adults

Other 2016 studies: determine density and % parasitism of BMSB in soybean and field corn (unknown host crop in GA); monitoring NE including *T. japonicus*


Glynn Tillman, USDA/ARS and  
David Buntin and Mike Toews, UGA

# Distribution of BMSB in SC



# BMSB in South Carolina Peach - 2016

- ❖ Pyramid traps baited with AgBio combo lure deployed at 6 sites in Spartanburg, SC
- ❖ Captured overwintering adults in spring of 2016
- ❖ Higher trap capture in peach in upstate SC than in peach in Coastal Plain in GA



Andy Rollins (Clemson University) and Ted Cottrell (USDA-ARS)

# **Alabama BMSB Team**

Rammohan Balusu

Savannah Duke

Henry Fadamiro

Kathy Flanders

Alana Jacobson

David Held

Xing Ping Hu


Ayanava Majumdar

Charles Ray

Tim Reed

Ron Smith

# Current Distribution Map



- Nymphal populations on soybean in various locations in north Alabama and Autauga county
- Damaging populations on soybean in Madison county and cotton in Autauga county
- Increase in stink bug damage on corn ears in north AL
- Serious nuisance pest in homes (more reports each year)

Map update: Charles Ray, Auburn University

# Projects Starting in 2016

## Fadamiro and Balusu Lab

- Evaluate the effect of trap color (yellow vs black) on capture of BMSB in vegetable and fruit crops (collaborative study with Ted Cottrell); 3 locations in AL- Fairhope, Clanton, and Auburn University (no BMSB captured yet); 1 location in GA - Byron
- Identify plant-based semiochemical attractants for BMSB

# Projects Starting in 2016

Smith, Reed, Duke, and Flanders

- Validating Treatment Thresholds and Determining Border Effect of Brown Marmorated Stink Bugs in Cotton

# Ongoing Projects

## Duke and Flanders

- Determine distribution of BMSB in AL
- Survey corn in North AL for ear damage

# Tennessee BMSB Team


Scott Stewart

Frank Hale

Karen Vail

Jerome Grant

# Brown Marmorated Stink Bug Distribution 2008-2015



# Pest Status in Tennessee


- BMSB expanding its range in TN
- BMSB is a significant nuisance/home pest in the eastern part of the state, especially in the urban areas of Knoxville and Nashville (to a lesser extent)
- Immatures and adults are damaging some fields of corn and soybean in the eastern 1/3 of the state
- Nuisance pest in the middle part of the state; suspect gaining a foothold and more common than known in crops in that part of the state.
- Low, but reproducing populations, in soybean near Memphis in 2015, but other than that, there have not been any confirmations of reproducing pest populations in the western part of the state.

Scott Stewart, University of Tennessee

## Planned Projects

- Monitoring in state
- Monitoring parasitism of BMSB in soybean

# Intensity of Brown Marmorated Stink Bug Populations in NC



- Widespread agricultural and nuisance pest.
- Locally intense agricultural and nuisance pest.
- Local hotspots in residential areas.
- Low level populations in isolated areas.
- No confirmed detections.

# NC Crops with Reproducing BMSB Populations and Economic Injury

- Apples
- Peaches
- Asian Pears
- Fruiting Vegetables (Econ. damage organic only)
- Corn
- Soybean (Damage not of econ. significance)
- Pecan
- Cotton



**Apple**



**Corn**



**Peach**



**Pear**



**Pecan**

Every nut on tree had this damage



**Pepper**




**Tomato**




**BMSB damage to various specialty crops**

Reevaluate economic threshold




Dominic Reising, NCSU

# BMSB in NC Cotton



## Cotton Production (480-lb Bales)



**Union & Stanly Counties :**  
only major cotton producing  
areas where BMSB is a  
serious pest.

These counties are one of the few places in the country where significant cotton acreage and high BMSB populations occur concomitantly. Mike Toews' grad student Whitney Hadden is sampling here in a corn-cotton-soybean agroecosystem.

# BMSB Monitoring in NC

- Prior to 2016, monitoring has been limited to western NC (mountains, western piedmont) where populations are most intense.
- 2016, Dominic Reisig and Mohammad-Amir Aghaee have established extensive monitoring program in cooperation with county agents in eastern NC

# Important Non-Crop Habitats

- Tree of Heaven
  - Catalpa
  - Paulownia
- Black cherry
- Wild grape
- Black walnut
  - Dogwood
  - Locust
- Yellowwood

# BMSB Natural Enemies

- Predation (generalist predators) more important than parasitism in cropping systems
- Most prevalent egg parasitoids
  - *Telenomus podsi* (rarely complete development in BMSB eggs)
  - *Anastatus mirabilis* and *A. redivii* (primarily in wooded habitats)
  - *Trissolcus brochymenae*, *T. edessae*, *T. euschisti*

# BMSB Research Efforts in NC

Jim Walgenbach, George Kennedy  
Dominic Reisig

- Overwintering ecology, phenology, and survivorship in different NC ecoregions.
- Management on tree fruits and fruiting vegetables in western NC.
- Impact of native natural enemies in managed and non-managed habitats. Monitoring for native species and potentially *T. japonicus*.
- Pheromone trapping studies this year are primarily looking at working with more user-friendly traps – small modified pyramid type traps and sticky traps.