


Factors that potentially mediate the ecological host range of *Trissolcus japonicus*

Paul S. Botch & Ernest S. Delfosse
Department of Entomology
Michigan State University

Brown Marmorated Stink Bug (BMSB)
BMSB Integrated Pest Management (IPM) Working Group Meeting
November 30, 2016


Trissolcus japonicus (Ashmead) (Hymenoptera: Platygasteridae)



High parasitism rates in the native range (60 to 80%)

Physiological Host-Specificity Tests (PHST)

Risk = Hazard x Exposure



PHST poses the risk hazard question,

“Can *T. japonicus* attack non-target species?”


To determine this,

- 23 Pentatomoidea spp. were tested; and
- Development and emergence occurred on 15 species in 11 genera.
- Therefore, the hazard prediction is that *T. japonicus* can complete development on some native Pentatomoidea.
- But, what about the exposure analysis?

Ecological Sieves

“Will *T. japonicus* attack non-target species?”


Physiological (Potential) Host Range



Sampling in Michigan

June – Sept, 2014 - 2016

- **Quantitative sampling (100 m transects sweeping/beating) in 4 Natural Areas:**
 - ❖ **5 Woodlots**
 - ❖ **8 Grasslands**
- **Qualitative sampling for BMSB.**



Current Distribution in Michigan Reported in 46 Counties

- Search by Date
- Search by Species
- Search by Geography
- Search by Project
- Search by Contributor


Reported Species Observations

Brown marmorated stink bug

* Please contact us for Phragmites data.



Messages: Please note that User Login is required to view contributor details.



BMSB Sampling

Host Plants:

June – Sept:

- Tree of Heaven
- Honeysuckle
- Ash
- Boxelder
- Eastern White Cedar
- Black Locust
- Pokeweed


Sept:

- Soybean




Pentatomidae in Woodlot Communities

- 15 species.
- *Banasa dimidiata* - 24%
- *Euschistus tristigmus luridus* - 24%



Pentatomoidea in Grassland Communities

- 16 species.
- *Cormelaena* spp.- 59%
- *Euschistus variolarious* - 16%



Adventive *T. japonicus* in the Field

Locations Where *T. japonicus* Have Been Found

As of 2015



Trissolcus japonicus (Ashmead) (Hymenoptera, Scelionidae) emerges in North America

Elijah J. Talamas¹, Megan V. Herlihy², Christine Dieckhoff^{3,4}, Kim A. Hoelmer⁴,
Matthew L. Buffington¹, Marie-Claude Bon⁵, Donald C. Weber²

¹ Systematic Entomology Laboratory, USDA/ARS c/o NMNH, Smithsonian Institution, Washington DC, USA ² Invasive Insect Biocontrol and Behavior Laboratory, USDA/ARS, BARC-West Beltsville MD, USA ³ Department of Entomology and Wildlife Ecology, University of Delaware, Newark, DE, USA ⁴ Beneficial Insects Introduction Research Unit, USDA/ARS, Newark, DE, USA ⁵ European Biological Control Laboratory, USDA/ARS, Montpellier, France

Discovery of an Exotic Egg Parasitoid of the Brown Marmorated Stink Bug, *Halyomorpha halys* (Stål) in the Pacific Northwest

Author(s): Joshua M. Milnes , Nik G. Wiman , Elijah J. Talamas , Jay F. Brunner , Kim A. Hoelmer , Matthew L. Buffington and Elizabeth H. Beers

Source: Proceedings of the Entomological Society of Washington, 118(3):466-470.

Published By: Entomological Society of Washington

DOI: <http://dx.doi.org/10.4289/0013-8797.118.3.466>

URL: <http://www.bioone.org/doi/full/10.4289/0013-8797.118.3.466>

MI Sentinel Egg Surveys




Parasitoids by Host Species. 2015

Host		Parasitoids		BMSB Eggs		
Common Name	Family	Species	Family	No. Put Out	Attacked No.	%
American hornbeam or Ironwood, <i>Carpinus caroliniana</i>	Betulaceae	<i>Trissolcus cosmopeplae</i>	Platygastridae	84	8	0.10
Milkweed, <i>Asclepias albicans</i>	Apocynaceae	<i>Ooencyrtus sp.</i>	Encyrtidae	84	8	0.10
Milkweed, <i>Asclepias albicans</i>	Apocynaceae	<i>Trissolcus euschisti</i>	Platygastridae	28	3	0.11
Wild Grape, <i>Vitis vinifera</i>	Vitaceae	<i>Trissolcus cosmopeplae</i>	Platygastridae	28	6	0.21
Wild Grape, <i>Vitis vinifera</i>	Vitaceae	<i>Ooencyrtus sp.</i>	Encyrtidae	27	4	0.15
Hawthorn, <i>Crataegus rhipidophylla</i>	Rosaceae	<i>Ooencyrtus sp.</i>	Encyrtidae	56	6	0.11
American basswood, <i>Tilia americana</i>	Tiliaceae	<i>Trissolcus cosmopeplae</i>	Platygastridae	28	1	0.04
American basswood, <i>Tilia americana</i>	Tiliaceae	<i>Trissolcus euschisti</i>	Platygastridae	28	2	0.07
American basswood, <i>Tilia americana</i>	Tiliaceae	<i>Trissolcus brochymenae</i>	Platygastridae	20	1	0.05
Totals				383	39	0.10

Ecological Sieves

Physiological (Potential) Host Range



Habitat Complexity Experiments



46 x 46 x 46 cm



1.5 x 1.5 x 2 m H


Habitat Complexity Experiments

Test	No. reps	No. egg masses parasitized			X^2 statistic	p-value
		BMSB	<i>T.c. accerra</i>	<i>P. maculiventris</i>		
Paired-choice	21	20	15	----	4.29	0.0384
	21	18	----	12	4.2	0.0404



Ecological Sieves

Physiological (Potential) Host Range



Semiochemicals


Host location by *Trissolcus basalis*:

- **Egg kairomones** (Bin *et al.* 1993).
- **Adult cuticular hydrocarbons** (Colazza *et al.* 2007).
- **Defensive secretions** (Laumann *et al.* 2009).
- **Chemical footprints on leaves** (Colazza *et al.* 2009).
- **Feeding and oviposition damage** (Colazza *et al.* 2004).




Semiochemicals

- **Y-tube Olfactometer.**
- **Odor Sources.**
 - ❖ Eggs.
 - ❖ Adult BMSB.
 - ❖ Chemical footprints.
 - ❖ Feeding damage.
- **Under Continued Evaluation.**




Ecological Sieves

Physiological (Potential) Host Range



Multiple-Species Choice Tests


- One, 24-h-old mated, naïve female *T. japonicus* placed in middle of arena.
- Scored behaviors:
 - Encounter with eggs;
 - Inspection of eggs by circling and antennal drumming;
 - Egg rejection (abandoning); and
 - Egg acceptance (oviposition).
- After 24-h, Petri dishes moved to an environmental chamber at 25°C; 60-80% RH; 16:8 L:D.
- Wasp removed from the arena after 24-h.
- Egg masses held separately until wasp or nymph emergence.



150 x 15 mm Petri dish arenas.


Multiple-Species Choice Tests

No. replicates in which eggs were accepted (blue) or rejected (orange) ($n = 30$)




Multiple-Species Choice Tests

Total Emergence ($n = 38$)



Multiple-Species Choice Tests

Total Emergence ($n = 38$)



Development on Non-Target Hosts

- **Does specificity differ?**


- Do compounds on eggs train wasps for preference?
- Genetic inclination?

- Tumlinson *et al.* 1993. How parasitic wasps find their hosts. *Scientific American*. March: 100-106.




- **Does fecundity differ?**

- Effects of phenotypic variation.

- Arakawa *et al.* 2004. Effects of host species on body size, fecundity, and longevity of *Trissolcus mitsukurii* (Hymenoptera: Scelionidae), a solitary egg parasitoid of stink bugs. *Appl. Entomol. Zool.* 39:177-181.
- Abram *et al.* 2015. Size-induced phenotypic reaction norms in a parasitoid wasp: an examination of life-history and behavioural traits. *Biological Journal of the Linnean Society*. (In Press).





Multiple-Species Choice Tests




Multiple-Species Choice Tests


BMSB Reared



T.c. accerra Reared



P. maculiventris Reared



Conclusions and Future Work

1. Risk = Hazard x Exposure, = “Can-Do” x “Will Do.”
 - Hazard is the innate capacity of a biological control agent to attack a non-target species and is determined in PHST; and
 - Exposure is determined by ecological sieves that may mediate the hazard.
 - A high hazard and a low exposure can mean that a potential biological control agent is “safe” to release.
2. The hazard analysis following PHST shows that *T. japonicus* can attack at least 11 genera of native Pentatomoidea.
3. The exposure analysis shows that there are some ecological sieves (habitat partitioning and host egg characteristics) that can mediate the potential hazard.
4. In the absence of mediating ecological sieves, we feel that the potential host range of *T. japonicus* is too broad, and it should not be approved for release.
5. However, since adventive populations of *T. japonicus* have been found in eastern and western U.S., this is a moot point, and efforts to identify ecological sieves and evaluate damage to native Pentatomoidea should be the focus of future research.

Acknowledgements



**USDA-APHIS Farm Bill Grant:
No. 11-13-8130-024 CA**

**This material was made possible,
in part, by Cooperative Agreement
8130-0024-CA 2016 from the United
States Department of Agriculture's
Animal and Plant Health Inspection
Service (APHIS). It may not
necessarily express APHIS' views.**



Delfosse Classical Biological Control Lab

**MSU AgBioResearch
MSU Extension
MSU Department of Entomology**