


Rhode Island - Set up ProtekNet around perimeter of 1/3 acre blueberry field at URI. No netting on top!




Set up 56 traps on outside around perimeter and set up 4 sentinel traps inside blueberry planting. Red cup traps baited with Suzuki Trap. Caught first SWD outside on 7/6/16. Caught first inside on 7/18/16. Did not catch any males until 8/1/16.

OUTSID E	6/27/1	7/6/1	7/11/1	7/18/1	7/25/1	8/1/1	8/9/1	8/15/1	8/22/1
Females	0	1	9	7	17	48	79	34	47
Males	0	0	0	0	0	1	5	8	10
INSIDE									
Females	0	0	0	2	3	1	11	11	30
Males	0	0	0	0	0	0	6	1	23

Female & male SWD in red cup traps baited with Suzuki Trap OUTSIDE ProtekNet around perimeter



Female & male SWD in red cup traps baited with Suzuki Trap INSIDE blueberry patch



6/24/16 - First swd-infested raspberry in commercial planting in RI.

7/6/16 - First swd adult in trap outside ProtekNet around blueberry planting URI.

7/18/16 - First swd-infested blueberry picked - (10 days later found swd pupae.)

7/18/16 - First swd adult inside ProtekNet.

8/1/16 - First male swd outside of netting - 4 weeks after first female.

8/9/16 - First male swd inside ProtekNet - 3 weeks after first female.

"Historical data" Harvest berries and set up to rear SWD

2013 - URI blues swd-free until 7/25/16 harvest

2014 - URI blues swd-free until 7/22/16 harvest

2015 - URI blues swd-free until 7/20/16 harvest (first found in Collins)

2016 - URI blues swd-free until 7/18/16 harvest (first found in Collins)

Submitted by Heather Faubert

Vermont – No trapping occurred at the UVM Hort Farm in 2016. SWD issues were erratic around the state - reports from the field indicated up to 50% loss in fall raspberries in Huntington (NW) and hardly any loss in Rochester (Central). They were moderate down south in fall raspberries - virtually no reports of blueberry infestation this year. Concord grapes in Vern Grubinger's home planting that are usually loaded with fruit flies, were not infested.

Submitted by Ann Hazelrigg