

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
White Grubs			*****										looks like drought damage and may 'pull up like carpeting'; rodents and crows may tear at turf. (Soil Sample)
Ants			*****										Mounds in turf. (Visually look for mounds)
Billbugs							*****						Turf wilts but does not respond to watering. Sawdust-like material in thatch. Turf is easy to pull up. Attacks all turf, favors Kentucky Bluegrass. (Scout with soapy flush)
Chinch Bugs							*****						Wilted or browned areas, most severe in sunny, sandy areas. Often found in dry turf areas with thick thatch and sandy soil. (Visually inspect thatch: can or core float method to confirm)
Cutworms						*****							Burrows surrounded by brown patches. Green frass may be present. More often found on closely mown turf. (Soapy flush)
Hyperodes						*****							Yellow patches begin at edges of turf areas. Damaged areas may appear water-soaked. Usually two generations per year. Found in Annual Bluegrass (Soapy flush for adults; core float for larvae)
Sod Webworm				*****		*****							Discrete brown areas which coalesce later. Most common in sunny areas. May cause damage in late spring. Found in all turf, especially sunny, dry, or sloped areas. (Soapy flush for larvae, Visually inspect for adults at twilight)
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	